

兔抗 ASCC2 多克隆抗体

- 中文名称: 兔抗 ASCC2 多克隆抗体
- 英文名称: Anti-ASCC2 rabbit polyclonal antibody
- 别 名: p100; ASC1p100
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: ASCC2
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

	ASCC2 (activating signal cointegrator 1 complex subunit
	2), also known as ASC1P100, is a 757 amino acid protei
	n that is ubiquitously expressed and contains one CUE
	domain. The gene encoding ASCC2 maps to human chr
	omosome 22, which houses over 500 genes and is the
	second smallest human chromosome. Mutations in sever
Background:	al of the genes that map to chromosome 22 are involv
	ed in the development of Phelan-McDermid syndrome,
	Neurofibromatosis type 2, autism and schizophrenia. Ad
	ditionally, translocations between chromosomes 9 and 2
	2 may lead to the formation of the Philadelphia Chrom
	osome and the subsequent production of the novel fusi
	on protein BCR-Abl, a potent cell proliferation activator





	found in several types of leukemias.
Applications:	ELISA, WB
Name of antibody:	ASCC2
Immunogen:	Synthetic peptide of human ASCC2
Full name:	activating signal cointegrator 1 complex subunit 2
Synonyms :	p100; ASC1p100
SwissProt:	Q9H1I8
ELISA Recommended dilution:	2000-5000
WB Predicted band size:	86 kDa
WB Positive control:	A431, 293T and hela cells
WB Recommended dilution:	500-2000

