

兔抗 BBS10 多克隆抗体

- 中文名称: 兔抗 BBS10 多克隆抗体
- 英文名称: Anti-BBS10 rabbit polyclonal antibody
- 别 名: Bardet-Biedl syndrome 10; C12orf58
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: BBS10
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

	This gene is a member of the Bardet-Biedl syndrome (B
	BS) gene family. Bardet-Biedl syndrome is an autosomal
	recessive disorder characterized by progressive retinal d
	egeneration, obesity, polydactyly, renal malformation an
	d mental retardation. The proteins encoded by BBS gen
	e family members are structurally diverse and the simila
Background:	r phenotypes exhibited by mutations in BBS gene family
	members is likely due to their shared roles in cilia form
	ation and function. Many BBS proteins localize to the b
	asal bodies, ciliary axonemes, and pericentriolar regions
	of cells. BBS proteins may also be involved in intracellul
	ar trafficking via microtubule-related transport. The prot
	ein encoded by this gene is likely not a ciliary protein



全国订货电话 4008-723-722

	but rather has distant sequence homology to type II ch aperonins. As a molecular chaperone, this protein may a ffect the folding or stability of other ciliary or basal bod y proteins. Inhibition of this protein's expression impairs
	ciliogenesis in preadipocytes. Mutations in this gene cau se Bardet-Biedl syndrome type 10.
Applications:	ELISA, WB, IHC
Name of antibody:	BBS10
Immunogen:	Fusion protein of human BBS10
Full name:	Bardet-Biedl syndrome 10
Synonyms:	C12orf58
SwissProt:	Q8TAM1
ELISA Recommended dilution:	5000-10000
	11 P

ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	30-150
WB Predicted band size:	81 kDa
WB Positive control:	PC-3 , A172 and HEPG2 cell lysates
WB Recommended dilution:	200-1000





全国订货电话 4008-723-722

