

## 兔抗 CCL27 多克隆抗体

- 中文名称: 兔抗 CCL27 多克隆抗体
- 英文名称: Anti-CCL27 rabbit polyclonal antibody
- 别 名: C-C motif chemokine ligand 27; ALP; ILC; CTAK; CTACK; PESKY; ESKINE; SCYA27
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗 原: CCL27
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

## 技术规格

Background:	This gene is one of several CC cytokine genes cluster ed on the p-arm of chromosome 9. Cytokines are a f amily of secreted proteins involved in immunoregulato ry and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin- associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine r eceptor 10 (CCR10). Studies of a similar murine protei n indicate that these protein-receptor interactions hav e a pivotal role in T cell-mediated skin inflammation.
Applications:	ELISA, IHC



## 全国订货电话 4008-723-722

Name of antibody:	CCL27
Immunogen:	Synthetic peptide of human CCL27
Full name:	C-C motif chemokine ligand 27
Synonyms:	ALP; ILC; CTAK; CTACK; PESKY; ESKINE; SCYA27
SwissProt:	Q9Y4X3
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	25-100

