

小鼠抗 FZD7 单克隆抗体

中文名称： 小鼠抗 FZD7 单克隆抗体

英文名称： Anti-FZD7 mouse monoclonal antibody

相关类别： 一抗

储 存： 冷冻 (-20°C)

宿 主： Mouse

抗 原： FZD7

反应种属： Human

标记物： Unconjugate

克隆类型： mouse monoclonal

技术规格

| | |
|--------------------------|--|
| Background: | Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD7 protein contains an N-terminal signal sequence, 10 cysteine residues typical of the cysteine-rich extracellular domain of Fz family members, 7 putative transmembrane domains, and an intracellular C-terminal tail with a PDZ domain-binding motif. FZD7 gene expression may downregulate APC function and enhance beta-catenin-mediated signals in poorly differentiated human esophageal carcinomas. |
| Applications: | ELISA, WB |
| Name of antibody: | FZD7 |
| Immunogen: | Fusion protein of human FZD7 |
| Full name: | frizzled class receptor 7 |
| Synonyms: | FzE3 |
| SwissProt: | O75084 |

| | |
|--------------------------|-------------------------|
| WB Predicted band size: | 64 kDa |
| WB Positive control: | HepG2 whole cell lysate |
| WB Recommended dilution: | 500-2000 |

