

DOK3 抗原（重组蛋白）

中文名称：DOK3 抗原（重组蛋白）

英文名称：DOK3 Antigen (Recombinant Protein)

储 存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to C terminal 300 amino acids of human DOK3

技术规格

Full name:	docking protein 3
Synonyms:	DOKL
Swissprot:	Q7L591
Gene Accession:	BC004564
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	DOK3 gene maps to chromosome 5q35.3. Dok3 was tyrosine phosphorylated by Src family members Lck, Fyn, and Lyn. Immunoprecipitation studies showed that Dok3 bound inhibitors SHIP and Csk but did not bind RasGAP. Dok3 binding to SHIP occurred via the SH2 domain. Dok3 also bound Csk via the Csk SH2 domain with possible involvement of the Csk SH3 domain as well. DOK proteins are enzymatically inert adaptor or scaffolding proteins. They provide a docking platform for the assembly of multimolecular signaling complexes. DOK3 is a negative regulator of JNK signaling in B-cells through interaction with INPP5D/SHIP1. May modulate ABL1 function

