

小鼠抗 STN1 单克隆抗体

中文名称： 小鼠抗 STN1 单克隆抗体

英文名称： Anti-STN1 mouse monoclonal antibody

别 名： STN1 subunit of CST complex; AAF44;OBFC1; AAF-44; RPA-32; bA541N10.2

相关类别： 一抗

储 存： 冷冻 (-20°C) 避光

宿 主： Mouse

抗 原： STN1

反应种属： Human

标 记 物： Unconjugate

克隆类型： mouse monoclonal

技术规格

Background:	OBFC1 and C17ORF68 (MIM 613129) are subunits of an alpha accessory factor (AAF) that stimulates the activity of DNA polymerase-alpha-primase (see MIM 176636), the enzyme that initiates DNA replication (Casteel et al., 2009 [PubMed 19119139]). OBFC1 also appears to function in a telomere-associated complex with C17ORF68 and TEN1 (C17ORF106; MIM 613130) (Miyake et al., 2009 [PubMed 19854130]).
Applications:	WB, IF, FC
Name of antibody:	STN1
Immunogen:	Fusion protein of human STN1
Full name:	STN1 subunit of CST complex
Synonyms:	AAF44; OBFC1; AAF-44; RPA-32; bA541N10.2
SwissProt:	Q9H668

WB Predicted band size:	42 kDa
WB Positive control:	HepG2, A549, Jurkat, MCF-7 cell lysates
WB Recommended dilution:	200-1000
IF positive control:	COS7 cells transiently transfected by pCMV6-ENTRY
IF Recommend dilution:	100-200
FC Recommend dilution:	50-100