

全国订货电话 4008-723-722

BRSK1 抗原(重组蛋白)

- 中文名称: BRSK1 抗原 (重组蛋白)
- 英文名称: BRSK1 Antigen (Recombinant Protein)
- 别 名: BR serine/threonine kinase 1; hSAD1
- 相关类别: 抗原
- 储 存: 冷冻 (-20℃)

概述

Fusion protein corresponding to a region derived from 579-778 amino acids of human BRSK1

技术规格

| Full name: | BR serine/threonine kinase 1 |
|--------------------|--|
| Synonyms: | hSAD1 |
| Swissprot: | Q8TDC3 |
| Gene Accession: | BC016681 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | The phosphorylation and dephosphorylation of proteins on serine a nd threonine residues is an essential means of regulating a broad r ange of cellular functions in eukaryotes, including cell division, hom eostasis and apoptosis. A group of proteins that are intimately invo lved in this process are the serine/threonine (Ser/Thr) protein kinas es. BRSK1 (BR serine/threonine-protein kinase 1), also known as SA D1, is a 794 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one UBA domain and one protein kinase domain. Expressed in a variety of tissues with highest expres sion in testis and brain, BRSK1 uses magensium as a cofactor to ca |



全国订货电话 4008-723-722

talyze the ATP-dependent phosphorylation of target proteins, includ ing Wee 1 and Cdc25B. Via its kinase activity toward proteins that are involved in microtubule assembly, BRSK1 plays an essential role in neuronal polarization and may be involved in regulating cell cycl e arrest in response to DNA damage, Two isoforms of BRSK1 exist due to alternative splicing events.