

兔抗 CYP2A13 多克隆抗体

中文名称: 兔抗 CYP2A13 多克隆抗体

英文名称: Anti-CYP2A13 rabbit polyclonal antibody

别名: cytochrome P450 family 2 subfamily A member 13 CPAD; CYP2A; CYP11A13

相关类别: 一抗

储存: 冷冻 (-20℃)

宿主: Rabbit

抗原: CYP2A13

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. Although its endogenous substrate has not been determined, it is known to metabolize 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, a major nitrosamine specific to tobacco. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q.
Applications:	ELISA, WB

Name of antibody:	CYP2A13
Immunogen:	Synthetic peptide of human CYP2A13
Full name:	cytochrome P450 family 2 subfamily A member 13
Synonyms:	CPAD; CYP2A; CYP2A13
SwissProt:	Q16696
ELISA Recommended dilution:	5000-10000
WB Predicted band size:	57 kDa
WB Positive control:	A549 cell lysates
WB Recommended dilution:	200-1000



