

STAT5A 抗原（重组蛋白）

中文名称：STAT5A 抗原（重组蛋白）

英文名称：STAT5A Antigen (Recombinant Protein)

储 存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to a region derived from 472-794 amino acids of human STAT5A

技术规格

| | |
|---------------------------|--|
| Full name: | Signal transducer and activator of transcription 5A |
| Synonyms: | MGF, STAT5 |
| Swissprot: | P42229 |
| Gene Accession: | BC027036 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. |

