

全国订货电话 4008-723-722

ILF3 抗原(重组蛋白)

中文名称: ILF3 抗原(重组蛋白)

英文名称: ILF3 Antigen (Recombinant Protein)

别 名: CBTF; DRBF; MMP4; MPP4; NF90; NFAR; NF110; NF90a; NF90b; NFAR2; TCP80; DRBP76; NF110b; NFAR-1; TCP110; MPHOSPH4; NF-AT-90

储存: 冷冻(-20℃)

相关类别: 抗原

概述:

Fusion protein corresponding to a region derived from 1-200 amino acids of human ILF3

技术规格:

Full name:	interleukin enhancer binding factor 3
Synonyms:	CBTF; DRBF; MMP4; MPP4; NF90; NFAR; NF110; NF90a; NF90b; NFA
	R2; TCP80; DRBP76; NF110b; NFAR-1; TCP110; MPHOSPH4; NF-AT-9
	0
Swissprot:	Q12906
Gene Accession:	BC064836
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	This gene encodes a double-stranded RNA (dsRNA) binding protein
	that complexes with other proteins, dsRNAs, small noncoding RNAs
	, and mRNAs to regulate gene expression and stabilize mRNAs. Thi
	s protein (NF90, ILF3) forms a heterodimer with a 45 kDa transcrip
	tion factor (NF45, ILF2) required for T-cell expression of interleukin
	2. This complex has been shown to affect the redistribution of nucl
	ear mRNA to the cytoplasm. Knockdown of NF45 or NF90 protein
	retards cell growth, possibly by inhibition of mRNA stabilization. In



全国订货电话 4008-723-722

contrast, an isoform (NF110) of this gene that is predominantly res tricted to the nucleus has only minor effects on cell growth when i ts levels are reduced. Alternative splicing results in multiple transcri pt variants encoding distinct isoforms.