

兔抗 PTPN6(Ab-564) 多克隆抗体

- 中文名称: 兔抗 PTPN6(Ab-564) 多克隆抗体
- 英文名称: Anti-PTPN6(Ab-564) rabbit polyclonal antibody
- 别名: MSK1; RLPK; MSPK1
- 相关类别: 一抗
- 储存: 冷冻(-20℃) 避光
- 宿 主: Rabbit
- 抗原: PTPN6(Ab-564)
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

| Background: | Serine/threonine-protein kinase that is required f or the mitogen or stress-induced phosphorylatio n of the transcription factors CREB1 and ATF1 a nd for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contribute s to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes. Phosphorylates CREB1 and ATF1 in respo nse to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and an isomycin. Plays an essential role in the control o f RELA transcriptional activity in response to TNF |
|-------------|---|
|-------------|---|



WB Recommended dilution:

全国订货电话 4008-723-722

| | and upon alucacarticaid accasiates in the extern |
|-------------------------|--|
| | and upon glucocorticoid, associates in the cytopl asm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression. In skeletal myobla sts is required for phosphorylation of RELA at 'S er-276' during oxidative stress. In erythropoietin- stimulated cells, is necessary for the 'Ser-727' ph osphorylation of STAT3 and regulation of its tra nscriptional potential. Phosphorylates ETV1/ER81 at 'Ser-191' and 'Ser-216', and thereby regulates its ability to stimulate transcription, which may b e important during development and breast tum or formation. Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A. Phosp horylates 'Ser-10' of histone H3 in response to mitogenics, stress stimuli and EGF, which results in the transcriptional activation of several immed iate early genes, including proto-oncogenes c-fo s/FOS and c-jun/JUN. May also phosphorylate 'S er-28' of histone H3. Mediates the mitogen- and stress-induced phosphorylation of high mobility group protein 1 (HMGN1/HMG14). In lipopolysac charide-stimulated primary macrophages, acts do wnstream of the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines. Fu nctions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inf lammatory cytokine interleukin 10 (IL10), via CRE B1 and ATF1 transcription factors. Plays a role in |
| | neuronal cell death by mediating the downstrea m effects of excitotoxic injury. |
| Applications: | WB |
| Name of antibody: | PTPN6(Ab-564) |
| Immunogen: | Synthesized peptide derived from C-terminal of human SHP-1. |
| Full name: | ribosomal protein S6 kinase, 90kDa, polypeptide 5 |
| Synonyms : | MSK1; RLPK; MSPK1 |
| SwissProt: | 075582 |
| WB Predicted band size: | 90 kDa |
| WB Positive control: | Jurkat cells lysate |
| | |

500-3000



全国订货电话 4008-723-722

