

全国订货电话 4008-723-722

## 兔抗 NCF4 多克隆抗体

- 中文名称: 兔抗 NCF4 多克隆抗体
- 英文名称: Anti-NCF4 rabbit polyclonal antibody
- 别 名: MGC3810; NCF; P40PHOX; SH3PXD4
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃) 避光
- 宿 主: Rabbit
- 抗 原: NCF4
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

## 技术规格

Background: ith the small G protein RAC1 and translocates to the membrane upon cell stimulation. This complex then activates flave cytochrome b, the membrane-integrated catalytic core of the e enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests it role in PI(3) kinase-mediated signaling events. The phospho	Background:	The protein encoded by this gene is a cytosolic regulatory c omponent of the superoxide-producing phagocyte NADPH-o xidase, a multicomponent enzyme system important for host defense. This protein is preferentially expressed in cells of m yeloid lineage. It interacts primarily with neutrophil cytosolic factor 2 (NCF2/p67-phox) to form a complex with neutrophil cytosolic factor 1 (NCF1/p47-phox), which further interacts w ith the small G protein RAC1 and translocates to the memb rane upon cell stimulation. This complex then activates flavo cytochrome b, the membrane-integrated catalytic core of th e enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests its role in PI(3) kinase-mediated signaling events. The phosphor ylation of this protein was found to negatively regulate the
--	-------------	---





	enzyme activity. Alternatively spliced transcript variants enco ding distinct isoforms have been observed.
Applications:	WB
Name of antibody:	NCF4
Immunogen:	Fusion protein of human NCF4
Full name:	neutrophil cytosolic factor 4, 40kDa
Synonyms :	MGC3810; NCF; P40PHOX; SH3PXD4
SwissProt:	Q15080
WB Predicted band size:	39 kDa
WB Positive control:	K562 cells
WB Recommended dilution:	500-2000

