

全国订货电话 4008-723-722

兔抗 PLAU 多克隆抗体

- 中文名称: 兔抗 PLAU 多克隆抗体
- 英文名称: Anti-PLAU rabbit polyclonal antibody
- 别名: ATF; QPD; UPA; URK; u-PA; BDPLT5
- 抗原: PLAU
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a serine protease involved in degrada tion of the extracellular matrix and possibly tumor cell m igration and proliferation. A specific polymorphism in thi s gene may be associated with late-onset Alzheimer's dis ease and also with decreased affinity for fibrin-binding. T his protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two -chain derivative in which a single disulfide bond connec ts the amino-terminal A-chain to the catalytically active, carboxy-terminal B-chain. This two-chain derivative is als o called HMW-uPA (high molecular weight uPA). HMW-u PA can be further processed into LMW-uPA (low molecul ar weight uPA) by cleavage of chain A into a short chai
-------------	--



全国订货电话 4008-723-722

	n A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA recep tor. Alternatively spliced transcript variants encoding diffe rent isoforms have been found for this gene.
Applications:	ELISA, IHC
Name of antibody:	PLAU
Immunogen:	Fusion protein of human PLAU
Full name:	plasminogen activator, urokinase
Synonyms :	ATF; QPD; UPA; URK; u-PA; BDPLT5
SwissProt:	P00749
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human gastric cancer and Human brain
IHC Recommend dilution:	10-50

