

兔抗 PPIG 多克隆抗体

- 中文名称: 兔抗 PPIG 多克隆抗体
- 英文名称: Anti-PPIG rabbit polyclonal antibody
- 别 名: CYP; SRCyp; SCAF10; CARS-Cyp
- 抗 原: PPIG
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 反应种属: Human, Mouse, Rat
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

| Background: | PPIases accelerate the folding of proteins. It catalyzes th e cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be implicated in the folding, trans port, and assembly of proteins. May play an important r ole in the regulation of pre-mRNA splicing. |
|-----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Applications: | ELISA, IHC |
| Name of antibody: | PPIG |
| Immunogen: | Fusion protein of human PPIG |
| Full name: | peptidylprolyl isomerase G |
| Synonyms: | CYP; SRCyp; SCAF10; CARS-Cyp |
| SwissProt: | Q13427 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human lung cancer |



IHC Recommend dilution:

35-200

