

兔抗

PRKAA1/PRKAA2(Phospho-Thr183/Thr172)多 克隆抗体

- 中文名称: 兔抗 PRKAA1/PRKAA2(Phospho-Thr183/Thr172)多克隆抗体
- 英文名称: Anti-PRKAA1/PRKAA2(Phospho-Thr183/Thr172) rabbit polyclonal antibody
- 别 名: AMPK; AMPKa1
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃) 避光
- 宿 主: Rabbit
- 抗 原: PRKAA1/PRKAA2(Phospho-Thr183/Thr172)
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

	The protein encoded by this gene is a catalytic subunit of
	the AMP-activated protein kinase (AMPK). AMPK is a hete
	rotrimer consisting of an alpha catalytic subunit, and non-
	catalytic beta and gamma subunits. AMPK is an important
Background:	energy-sensing enzyme that monitors cellular energy statu
	s. In response to cellular metabolic stresses, AMPK is activ
	ated, and thus phosphorylates and inactivates acetyl-CoA
	carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-C
	oA reductase (HMGCR), key enzymes involved in regulatin



全国订货电话 4008-723-722

	g de novo biosynthesis of fatty acid and cholesterol. Stud ies of the mouse counterpart suggest that this catalytic s ubunit may control whole-body insulin sensitivity and is n ecessary for maintaining myocardial energy homeostasis d uring ischemia.
Applications:	WB, IHC
Name of antibody:	PRKAA1/PRKAA2(Phospho-Thr183/Thr172)
Immunogen:	Peptide sequence around phosphorylation site of threonin e 183/threonine172(L-R-T(p)-S-C) derived from Human AM $PK\alpha 1/AMPK\alpha 2$.
Full name:	protein kinase AMP-activated catalytic subunit alpha 1/ pr otein kinase AMP-activated catalytic subunit alpha 2
Synonyms:	AMPK; AMPKa1; AMPK2; PRKAA; AMPKa2
SwissProt:	Q13131/P54646
IHC positive control:	Human lung carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	64 kDa
WB Positive control:	C2C12 cells lysates treated with serum or calf intestinal p hosphatase(CIP)
WB Recommended dilution:	500-1000





